



## *Acrocephalus dumetorum*

---

<b>Annex I</b>	No
<b>International action plan</b>	No

Blyth's Reed-warbler, *Acrocephalus dumetorum*, is a species of passerine bird in the warbler family found in wetland and woodland and forest ecosystems. It is a widespread summer visitor to central Russia and adjoining areas of northern and central Europe (Bird in Europe 2004). This species has adapted to varied habitats and is not necessarily found close to water; it is more arboreal than most members of the genus. It breeds in a wide range of shrubby growth with tall herbage, young forests, overgrown clearings, birch (*Betula*) and willow (*Salix*) scrub in floodland groves, forest steppes, bush-encroached farmland and swamp and other waterside margins (European Red List 2015).

*Acrocephalus dumetorum* has a breeding population size of 76500-180000 pairs and a breeding range size of 464000 square kilometres in the EU27. The breeding population trend in the EU27 is Increasing in the short term and Increasing in the long term.


The EU population status of *Acrocephalus dumetorum* was assessed as Secure, because the species does not meet any of the IUCN Red List criteria for threatened or Near Threatened, or the criteria for Depleted or Declining (the EU27 population or range has not declined by 20% or more since 1980).

## Assessment of status at the European level

Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend		Population status
	Short term	Long term		Short term	Long term		Short term	Long term	
76500 - 180000 p	+	+	464000	+	+				Secure

See the endnotes for more information<sup>i</sup>

***Acrocephalus dumetorum***  
Report under the Article 12 of the Birds Directive





***Acrocephalus dumetorum***  
Report under the Article 12 of the Birds Directive

## Trends at the Member State level

MS/Ter.	% in EU27	Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend	
			Short term	Long term		Short term	Long term		Short term	Long term
EE	16.7	60000 - 120000 p	+	+	45700	+	+			
FI	51.6	16000 - 59000 p	+	+	227700	x	+			
LT	2.9	500 - 700 p	+	+	22500	+	+			
LV	13.7		x	x	61217	x	x			
PL		6 - 28 cmales	x	x		x	x			
SE	15.1	50 - 200 p	+	+	107200	+	x			

See the endnotes for more information<sup>ii</sup>

***Acrocephalus dumetorum***  
 Report under the Article 12 of the Birds Directive



***Acrocephalus dumetorum***  
*Report under the Article 12 of the Birds Directive*

*Short-term winter population trend was not reported for this species.*

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at:  
<http://bd.eionet.europa.eu/article12/summary?period=1&subject=A679> .

***Acrocephalus dumetorum***  
*Report under the Article 12 of the Birds Directive*

<sup>i</sup> **Assessment of status at the European level:** The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

<sup>ii</sup> **Species trends at the Member State level:** The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.