European Environment Agency *European Topic Centre on Biological Diversity*



Acrocephalus scirpaceus

Annex I	No
International action plan	No

Eurasian Reed-warbler, *Acrocephalus scirpaceus*, is a species of passerine bird in the warbler family found in wetland and river and lake ecosystems. It is broadly distributed across the whole of Europe, with somewhat patchy distribution across the Southern and South-Eastern Europe. This species breeds mainly in mature beds of reed (*Phragmites*) on the shores of lakes and fish ponds, and along rivers and ditches and locally, breeds in willow bushes in marshland, in reeds on edges of brackish lakes, exceptionally in corn fields. It also forages in adjacent herbaceous vegetation, scrub and low trees, such as willows (*Salix*) (European Red List 2015).

Acrocephalus scirpaceus has a breeding population size of 1490000-2660000 pairs and a breeding range size of 2470000 square kilometres in the EU27. The breeding population trend in the EU27 is Stable in the short term and Stable in the long term.

The EU population status of *Acrocephalus scirpaceus* was assessed as Secure, because the species does not meet any of the IUCN Red List criteria for threatened or Near Threatened, or the criteria for Depleted or Declining (the EU27 population or range has not declined by 20% or more since 1980).

Acrocephalus scirpaceus


Report under the Article 12 of the Birds Directive

Assessment of status at the European level

Breeding population size	Breeding population trend		Range	Breeding range trend		Winter	Winter population trend		Population
	Short term	Long term	area	Short term	Long term	population size	Short term	Long term	status
1490000 - 2660000 p	0	0	2470000						Secure

See the endnotes for more informationⁱ

Acrocephalus scirpaceus Report under the Article 12 of the Birds Directive



Acrocephalus scirpaceus

Report under the Article 12 of the Birds Directive


Trends at the Member State level

MS/Ter.	% in	Breeding		ding on trend	Range		ng range end	Winter	Winter po tre	
W3/Ter.	EU27	population size	Short term	Long term	area	Short term	Long term	population size	Short term	Long term
AT	1.8	50000 - 65000 p	0	x	64964	0	0			
BE	1.9	10000 - 15000 p	0	+	27176	+	+			
BG	1.0	5000 - 10000 p	+	0	50100	+	0			
CY	0.1	100 - 300 p	x	x	5100	0	x			
CZ										
DE	21.3	110000 - 180000 p	F	+	324128	0	0			
DK	3.1	34000 - 34000 p	-	-	42793	0	0			
EE	1.5	10000 - 20000 p	0	0	52300	0	0			
ES	14.2	339500 - 660000 p	0	0	405970	0	0			
FI	2.8	20000 - 30000 p	-	-	93600	x	0			
FR	12.9	35000 - 70000 p	x		378100	X	-			
GR										
HU	1.4	70000 - 137000 p		x	93011	0	x			
IE	0.2	65 - 100 p	+	+	3400	+	+			
IT	7.4	30000 - 60000 p	-	-	221600	+	+			
LT	3.8	20000 - 40000 p	+	+	71300	0	0			
LU	0.1	200 - 250 p	+	+	2540	0	0			
LV	2.1		x	x	59910	x	0			
MT		5 - 8 p	0	+	3	+	+			
NL	2.9	156457 - 260762 p	+	0	42483	0	0			
PL		120000 - 190000 p	0	x		X	x			
PT	1.0	2500 - 10000 p	-	0	36300	0	0			
RO	0.6	127868 - 161130 p	x	x	142500	x	x			
SE	10.6	174000 - 405000 p	0	-	217600	0	x			
SI	0.2	200 - 300 p	0	+	2702	x	+			
SK	0.9	10000 - 20000 p	0	0	24098	0	0			
UK	8.0	100000 - 160000 p	+	+	109700	+	+			

See the endnotes for more informationⁱⁱ

Acrocephalus scirpaceus

Report under the Article 12 of the Birds Directive



Short-term winter population trend was not reported for this species.

Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article_12/reference_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats reported as 'high', for some species there were less than ten pressures and threats reported as highly important.

Ten most frequently reported 'highly important' pressures and threats

No 'highly important' pressures or threats were reported.

Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
AT	breeding	NO	
BE	breeding	NO	
BG	breeding	NO	
CY	breeding	NO	
DE	breeding	NO	
DK	breeding	NO	
EE	breeding	NO	
ES	breeding	NO	
FI	breeding	NO	
FR	breeding	NO	
HU	breeding	NO	
IE	breeding	NO	
IT	breeding	NO	
LT	breeding	NO	
LU	breeding	YES	70

Report under the Article 12 of the Birds Directive

MS/territory	season	SPA trigger	% coverage
LV	breeding	NO	
MT	breeding	NO	
NL	breeding	NO	
PL	breeding	NO	
PT	breeding	NO	
RO	breeding	NO	
SE	breeding	NO	
SI	breeding	NO	
SK	breeding	NO	
UK	breeding	NO	

See the endnotes for more informationⁱⁱⁱ

Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
4.0	Other wetland-related measures	40
4.2	Restoring/improving the hydrological regime	20
6.1	Establish protected areas/sites	20
6.3	Legal protection of habitats and species	20

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at: http://bd.eionet.europa.eu/article12/summary?period=1&subject=A297. Report under the Article 12 of the Birds Directive

ⁱ Assessment of status at the European level: The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

ⁱⁱSpecies trends at the Member State level: The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

ⁱⁱⁱ**Percentage of coverage by Natura 2000 sites:** In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.