



Scolopax rusticola

Europe/South & West Europe & North Africa

Annex I	No
International action plan	No

Eurasian Woodcock, *Scolopax rusticola*, is a species of wader found in woodland and forest ecosystems.

Scolopax rusticola has a breeding population size of 728000-1480000 calling males and a breeding range size of 1890000 square kilometres in the EU27. The breeding population trend in the EU27 is Decreasing in the short term and Increasing in the long term.

The EU population status of *Scolopax rusticola* was assessed as Secure, because the species does not meet any of the IUCN Red List criteria for threatened or Near Threatened, or the criteria for Depleted or Declining (the EU27 population or range has not declined by 20% or more since 1980).



Scolopax rusticola [Europe/South & West Europe & North Africa]
Report under the Article 12 of the Birds Directive

Assessment of status at the European level

Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend		Population status
	Short term	Long term		Short term	Long term		Short term	Long term	
728000 - 1480000 cmales	-	+	1890000						Secure

See the endnotes for more informationⁱ

Scolopax rusticola [Europe/South & West Europe & North Africa]
Report under the Article 12 of the Birds Directive




Scolopax rusticola [Europe/South & West Europe & North Africa]
Report under the Article 12 of the Birds Directive

Trends at the Member State level

MS/Ter.	% in EU27	Breeding population size	Breeding population trend		Range area	Breeding range trend		Winter population size	Winter population trend	
			Short term	Long term		Short term	Long term		Short term	Long term
AT	2.1	2000 - 10000 cmales	x	x	70861	x	x			
BE	1.2	1000 - 5000 cmales	0	0	15394	-	0			
BG	0.4	90 - 180 cmales	x	x	15100	x	x			
CZ										
DE	22.0	20000 - 39000 cmales	0	0	305184	0	0		x	x
DK	1.0	2000 - cmales	0	+	21426	-	-			
EE	3.8	30000 - 60000 cmales	0	0	49400	0	0			
ES	1.9	3600 - 4000 cmales	0	0	49206	0	0	640000 - 680000 i	0	0
ESIC	0.2	1000 - 2500 cmales	x	x	1975	x	x			
FI	18.1	150000 - 220000 p	0	+	300600	x	+			
FR	5.2	10000 - 10000 cmales	0	0	186700	0	-			
GIB								1 - 5 i	0	0
GR										
HU	0.1	10 - cmales	x	x	14113	x	x			
IE	1.1		0	x	24700	0	-			
IT	1.9	50 - 150 cmales	x	x	51900	+	+			
LT	5.4	10000 - 20000 cmales	0	0	71000	0	0			
LU			x	x		x	x			
LV	4.4	10925 - 104363 cmales	-	+	64338	x	0			
NL	1.2	1000 - 5000 cmales	x	x	33568	-	0			
PL		20000 - 100000 cmales	x	x		x	x			
PTAC	0.6	1900 - 3800 cmales	0	x	6500	x	x			
PTMA	0.1	- 162 cmales	-	-	900	+	x			
RO		1000 - 5000 cmales	x	x	66400	x	x			
SE	19.7	396000 - 774000 cmales	0	0	421600	0	x			
SI	0.1	20 - 40 cmales	-	-	1361	-	-			
SK	2.1	1300 - 2500 cmales	-	-	36196	0	0			
UK	7.5	64000 - 100000 cmales	-	-	86600	-	-			

See the endnotes for more informationⁱⁱ

Scolopax rusticola [Europe/South & West Europe & North Africa]
 Report under the Article 12 of the Birds Directive



Scolopax rusticola [Europe/South & West Europe & North Africa]
 Report under the Article 12 of the Birds Directive



Scolopax rusticola
 Europe/South & West Europe & North Africa sub-population
 Short-term winter population trend

	Decreasing		Extinct		<= 3] No. of individuals
	Increasing		Present		<= 660 000	
	Stable or fluctuating		EU Member States		<= n .a.	
	Unknown trend		Outside data coverage		<= n .a.	

Main pressures and threats reported by Member States

For the bird species triggering SPA classification Member States were asked to report the 20 most important pressures and threats using an agreed hierarchical list which can be found on the Article 12 Reference Portal (http://bd.eionet.europa.eu/activities/Reporting/Article_12/reference_portal). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. The table below only contains information from Member States, where a species triggers SPA classification. Pressures and threats were ranked in three classes 'high, medium and low importance', the table below only shows pressures and threats classed as 'high', for some species there were less than ten pressures and threats reported as highly important.

Ten most frequently reported 'highly important' pressures and threats

Code	Activity	Frequency
F03	Hunting and collection of terrestrial wild animals	100

Proportion of population covered by the Natura 2000 network

For the bird species triggering SPA classification Member States were asked to report the size of a species population occurring within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size.

Percentage of coverage by Natura 2000 sites

MS/territory	season	SPA trigger	% coverage
AT	breeding	NO	
BE	breeding	NO	
BG	breeding	NO	
DE	breeding	NO	
DE	winter	NO	
DK	breeding	NO	
EE	breeding	NO	
ES	breeding	YES	1.03
ES	winter	YES	17.27
ESIC	breeding	NO	
FI	breeding	NO	
FR	breeding	NO	
GIB	winter	YES	100
HU	breeding	NO	
IE	breeding	NO	
IT	breeding	NO	
LT	breeding	NO	
LU	breeding	NO	

Scolopax rusticola [Europe/South & West Europe & North Africa]
Report under the Article 12 of the Birds Directive

MS/territory	season	SPA trigger	% coverage
LV	breeding	NO	
NL	breeding	NO	
PL	breeding	YES	x
PTAC	breeding	NO	
PTMA	breeding	NO	
RO	breeding	NO	
SE	breeding	NO	
SI	breeding	NO	
SK	breeding	NO	
UK	breeding	NO	

See the endnotes for more informationⁱⁱⁱ

Most frequently reported conservation measures

For the bird species triggering SPA classification Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 12 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.3	Legal protection of habitats and species	44
6.1	Establish protected areas/sites	33
7.1	Regulation/ Management of hunting and taking	22

This information is derived from the Member State national reports submitted to the European Commission under Article 12 of the Birds Directive in 2013 and covering the period 2008-2012. More detailed information, including the MS reports, is available at: <http://bd.eionet.europa.eu/article12/summary?period=1&subject=A155> .

Scolopax rusticola [Europe/South & West Europe & North Africa]

Report under the Article 12 of the Birds Directive

ⁱ **Assessment of status at the European level:** The EU assessments of birds population status was made by the European Red List of Birds Consortium (under contract with the European Commission)

The EU27 population trends were assessed using these categories: '+' Increasing, '0' Stable, 'F' Fluctuating, '-' Decreasing, 'xu' Uncertain and 'x' Unknown. The breeding population size is estimated in majority of the cases as 'p' number of pairs. Alternative population units used are: 'males' number of males, 'i' number of individuals, 'cmales' number of calling males and 'bfem' number of breeding females. The winter population size is estimated as number of individuals.

ⁱⁱ **Species trends at the Member State level:** The percentage of the EU27 species population occurring in the Member States (% in EU27) is calculated based on the population size reported by the Member States.

ⁱⁱⁱ **Percentage of coverage by Natura 2000 sites:** In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece and for non-Annex I species in the Czech Republic.