

A2.32 Polychaete/oligochaete-dominated upper estuarine Pontic littoral mud

Summary

The habitat is present in the Black Sea estuaries where mud is present. It is not present in the Sea of Marmara. Eutrophication is the main historic pressure on this habitat. Additional pressures include: coastal development and chemical pollution. Conservation and management measures relevant to this habitat include: measures to maintain physical and biological integrity, improvement of water quality controls, coastal development controls, survey and monitoring programmes, increased public awareness and designation of MPAs.

Synthesis

Detailed information on the abundance and extent of this habitat is lacking. Information on the quantity and quality of this habitat including historical or recent trends is unknown. For the purposes of Red List assessment this habitat is considered to be Data Deficient.

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Sub-habitat types that may require further examination

None

Habitat Type

Code and name

A2.32 Polychaete/oligochaete-dominated upper estuarine Pontic littoral mud

No characteristic photographs currently available for this habitat.

Habitat description

Upper estuarine sandy mud and mud shores, in areas with significant freshwater influence. Littoral mud typically forms mudflats, though dry compacted mud can form steep and even vertical structures, particularly at the top of the shore adjacent to saltmarshes. Little oxygen penetrates these cohesive sediments, and an anoxic layer is often present within millimetres of the sediment surface.

There are three oligochaete dominated upper estuarine mud biotopes variously dominated by polychaetes, oligochaetes and chironomids. The Pontic 'Camca' biotope which is associated with this habitat is unique to Danube River mouths.

Indicators of quality:

Both biotic and abiotic indicators have been used to describe marine habitat quality. These include: the presence of characteristic species and those which are sensitive to the pressures the habitat may face; water quality parameters; levels of exposure to particular pressure and more integrated indices which describe habitat structure and function, such as trophic index, or successional stages of development in habitats that have a natural cycle of change over time.

There are no commonly agreed indicators of quality for this habitat, although particular parameters may

have been set in certain situations e.g. protected features within Natura 2000 sites, where reference values have been determined and applied on a location-specific basis.

Characteristic species:

The upper estuarine mud communities support few infaunal species and are principally characterised by a restricted range of polychaetes and oligochaetes. Chironomids are present in the upper estuary where fresh water conditions are prevalent.

Classification

This habitat may be equivalent to, or broader than, or narrower than the habitats or ecosystems in the following typologies.

EUNIS (v1405):

Level 4. A sub-habitat of 'Pontic littoral mud' (A2.3).

Annex 1:

1130 Estuaries

1140 Mudflats & sandflats not covered at low tide

1160 Large shallow inlets and bays

MAES:

Marine - Marine inlets and transitional waters

Marine - Coastal

MSFD:

Littoral sediment

EUSEaMap:

Not mapped

IUCN:

9.10 Estuaries

12.4 Mud Shoreline and Intertidal Mud Flats

Does the habitat type present an outstanding example of typical characteristics of one or more biogeographic regions?

Unknown

Justification

There is insufficient knowledge and information on this habitat to state whether it is an outstanding example of this biogeographic region.


Geographic occurrence and trends

Region	Present or Presence Uncertain	Current area of habitat	Recent trend in quantity (last 50 yrs)	Recent trend in quality (last 50 yrs)
<i>Black Sea</i>	Black Sea: Present	Unknown Km ²	Unknown	Unknown

Extent of Occurrence, Area of Occupancy and habitat area

	Extent of Occurrence (EOO)	Area of Occupancy (AOO)	Current estimated Total Area	Comment
<i>EU 28</i>	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.
<i>EU 28+</i>	Unknown Km ²	Unknown	Unknown Km ²	The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO.

Distribution map



This habitat occurs in the Black Sea but there is insufficient data to produce a map of its distribution or calculate EOO and AOO.

How much of the current distribution of the habitat type lies within the EU 28?

It is unknown how much of this habitat is hosted by the EU 28 in the Black Sea.

Trends in quantity

There is insufficient data to accurately assess changes in quantity of the habitat.

- Average current trend in quantity (extent)

EU 28: Unknown

EU 28+: Unknown

- Does the habitat type have a small natural range following regression?

Unknown

Justification

The habitat is known to occur in the Black Sea but there is insufficient data to accurately calculate EOO and AOO. There is insufficient data to accurately assess whether the habitat has undergone a significant decline in the last 50 years.

- Does the habitat have a small natural range by reason of its intrinsically restricted area?

Unknown

Justification

There is insufficient data and knowledge on this habitat to state whether it has a small natural range by reason of an intrinsically restricted area.

Trends in quality

There is insufficient data to accurately assess changes in quality of the habitat.

- Average current trend in quality

EU 28: Unknown

EU 28+: Unknown

Pressures and threats

Eutrophication as a result of nutrient enrichment (N, P and organic matter) is the most significant historic pressure on the habitat. Hypoxic and anoxic events resulted in mass mortality of faunal species. Since the 1990s this pressure has reduced due to tighter controls on pollution in the catchment of the Danube and other rivers which enter the north-west Black Sea. Whilst this pressure is now reduced it is still a continuing threat in the current and future periods. This is especially true for non EU countries surrounding the Black Sea which are not bound by the agreements such as the Water Framework Directive (WFD).

Coastal developments including the construction of marinas and slipways, sediment extraction, the widening and dredging of channels, creation of artificial beaches, road construction and sea defences. These activities may alter the hydrological regime which will in turn affect the character and viability of the habitat.

Chemical pollution. This is a threat of current and future importance which at its most severe can result in species can lead to mortality. High mortality rates can lead to a reduction in extent. Lower mortality rates will result in a reduction in habitat quality. Chemical pollution may also affect the size and growth rate of some of the associated fauna.

List of pressures and threats

Urbanisation, residential and commercial development

Other urbanisation, industrial and similar activities

Pollution

Nutrient enrichment (N, P, organic matter)

Input of contaminants (synthetic substances, non-synthetic substances, radionuclides) - diffuse sources, point sources, acute events

Conservation and management

Conservation and management measures which would benefit this habitat include measures to maintain physical and biological integrity, including pollution control and regulation, water quality management, coastal development controls, contingency plans to be followed in the event of a major pollution incident, survey and monitoring programmes, raised public awareness of ecological value and vulnerability, enhanced legal protection for occurrences of the habitat and key species (e.g. additions to the EU Habitats Directive, the establishment of a unified list of Black Sea species and habitats requiring conservation measures) and the designation of MPAs.

List of conservation and management needs

Measures related to wetland, freshwater and coastal habitats

Restoring/Improving water quality

Measures related to marine habitats

Other marine-related measures

Measures related to spatial planning

Establish protected areas/sites

Legal protection of habitats and species

Measures related to urban areas, industry, energy and transport

Other measures

Conservation status

Annex 1:

1130 MBL5 U1

When severely damaged, does the habitat retain the capacity to recover its typical character and functionality?

There is insufficient data and knowledge on this habitat to state whether it has a small natural range by reason of an intrinsically restricted area.

Effort required

10 years
Unknown

Red List Assessment

Criterion A: Reduction in quantity

Criterion A	A1	A2a	A2b	A3
EU 28	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %

There is insufficient data on changes in quantity. This habitat has been assessed as Data Deficient under criterion A.

Criterion B: Restricted geographic distribution

Criterion B	B1				B2				B3
	EOO	a	b	c	AOO	a	b	c	
EU 28	unknown Km ²	Unknown							
EU 28+	unknown Km ²	Unknown							

This habitat occurs in the Black Sea but there is insufficient data to calculate EOO and AOO.

Criterion C and D: Reduction in abiotic and/or biotic quality

Criteria C/D	C/D1		C/D2		C/D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion C	C1		C2		C3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion D	D1		D2		D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%
EU 28+	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%

Experts consider there to be insufficient data to conduct an assessment using criteria C/D.

Criterion E: Quantitative analysis to evaluate risk of habitat collapse

Criterion E	Probability of collapse
EU 28	unknown
EU 28+	unknown

There is no quantitative analysis available to estimate the probability of collapse of this habitat type.

Overall assessment "Balance sheet" for EU 28 and EU 28+

	A1	A2a	A2b	A3	B1	B2	B3	C/D1	C/D2	C/D3	C1	C2	C3	D1	D2	D3	E
EU28	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD
EU28+	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD	DD

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Data Deficient	-	Data Deficient	-

Confidence in the assessment

Low (mainly based on uncertain or indirect information, inferred and suspected data values, and/or limited expert knowledge)

Assessors

S. Beal, G. Komakhidze, D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Contributors

S. Beal, G. Komakhidze, V. Kopy D. Micu, V. Mihneva, N. Milchakova, B. Yokes

Reviewers

S. Gubbay

Date of assessment

19/03/2015

Date of review

16/12/2015

References

Bacescu, M. C., Muller G. I., Gomoiu, M-T. 1971. . Cercetari de ecologie bentica in Marea Neagra (analiza cantitativa, calitativa si comparata a faunei bentice pontice). *Ecologie Marina* vol. IV. Editura Academiei R.S.R., Bucuresti, 357 pp.

Kiseleva, M. I. 1981. *Benthos of Black Sea mobile substrates*. Naukova dumka, Kiev, pp 165.

Konsulov, A. 1998. *Black Sea Biological Diversity: Bulgaria. Volume 5 of Black Sea environmental series*. United Nations Publications, New York, USA.

Marinov, T. 1990. *The zoobenthos from the Bulgarian Sector of the Black Sea*. Publishing house of the Bulgarian Academy of Sciences, Sofia, pp 195 (in Bulgarian).

Micu, D., Zaharia, T., Todorova, V., Niță, V. 2007. *Romanian Marine Habitats of European Interest*. Punct Ochit Publishers, Constanța, Romania.

Micu, D. 2008. Open Sea and Tidal Areas. In: Gafta D. and Mountford J.O. (eds.) *Natura 2000 Habitat Interpretation Manual for Romania*. EU publication no. EuropeAid/121260/D/SV/RO, 101pp. ISBN 978-973- 751-697-8.

Micu, D., Zaharia, T., Todorova, V. 2008. Natura 2000 habitat types from the Romanian Black Sea. In: Zaharia T, Micu D, Todorova V, Maximov V, Niță V. *The development of an indicative ecologically coherent network of marine protected areas in Romania*. Romart Design Publishing, Constanta, Romania.

Moncheva. S., Todorova, V., (eds). 2013. *Initial assessment of the marine environment*. Article 8, MSFD 2008/56/EC and NOOSMV (2010). 500p

Petranu, A. 1997. *Black Sea Biological Diversity: Romania. Volume 4 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.

Zaitsev, Y. P., Alexandrov, B. G. 1998. *Black Sea Biological Diversity: Ukraine. Volume 7 of the Black Sea Environmental Series*. United Nations Publications, New York, USA.