

A5.25 Atlantic upper circalittoral fine sand

Summary

This habitat consists of clean fine sands with less than 5% silt/clay in deeper water in tide-swept channels of marine inlets or on the open coast, extending offshore in depths of over 15-20 m. It is characterised by a wide range of echinoderms as well as polychaetes and bivalves. Frequently occurring species include the brittlestars *Amphiura filiformis*, *Ophiura albida* and *O. ophiura*, the anemone *Cerianthus lloydii*, and polychaete worms such as *Lanice conchilega*.

Disturbance of the substratum due to intensive fishing activities using bottom trawls or dredges may damage or modify infaunal communities. Smothering is not likely to cause considerable mortality to the associated species, however polluted sediments may lead to a change in the species composition by altering the sedimentary environment. Inshore dredging or construction works can also change the sediment structure and therefore community structure, by affecting the hydrodynamics or sedimentary processes. In addition, synthetic and non-synthetic compound contamination from dumped dredge spoil and local discharges could cause a decline in the species richness within this habitat. Beneficial management measures include control and restrictions on mobile demersal fishing, dredging and offshore construction works. The regulation of effluent discharges can also support the conservation of this habitat.

Synthesis

This habitat is relatively common and has a considerable natural range in the North East Atlantic although a patchy distribution.

Most sedimentary benthic systems on the continental shelf of Europe are believed to have been modified by fishing activities in the last 100 years, particularly by mobile demersal gears, and this habitat remains under fishing pressure. Data from a single year, 2013/2014, has revealed that than 40% of circalittoral fine sands and muddy sand were subject to trawling fishing pressure in the North Sea, with over 10% of this being interpreted a high or moderate pressure. When combining data for the North Sea and Celtic Sea more than 80% of this habitat type is considered to have been subject to such fishing pressure. Given that this is based on a single year of data, and that this type of pressure has been taking place for decades, it is likely to be an underestimate of the total area of this habitat which has been subject to such pressure. Disturbance of the substratum due to intensive fishing activities using bottom trawls or dredges can damage or modify infaunal communities, with burrowing echinoderms and bivalves being particularly vulnerable and therefore affect habitat quality.

Expert opinion is that there has been a very substantial reduction in quality of this habitat, most likely an intermediate decline affecting more than 80% of its extent although it is clear that in some locations there has also been a severe decline. The severity will depend on factors such as the intensity and frequency of disturbance. This habitat has therefore been assessed as Endangered for both the EU 28 and EU 28+ because of both past and likely continuing declines in quality.

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Endangered	C/D1	Endangered	C/D1

Sub-habitat types that may require further examination

None.

Habitat Type

Code and name

A5.25 Atlantic upper circalittoral fine sand

No characteristic photographs of this habitat currently available.

Habitat description

This habitat consists of clean fine sands with less than 5% silt/clay in deeper water in tide-swept channels of marine inlets or on the open coast, extending offshore in depths of over 15-20 m. Off the coast of Portugal it has been reported in depths of 20-37m in areas of moderately strong current. It is characterised by a wide range of echinoderms as well as polychaetes and bivalves. Frequently occurring species include the brittlestar *Amphiura filiformis*, *Ophiura albida* and *O. ophiura*, the anemone *Cerianthus lloydii*, and polychaete worms such as *Lanice conchilega*.

Indicators of quality:

Both biotic and abiotic indicators have been used to describe marine habitat quality. These include: the presence of characteristic species as well as those which are sensitive to the pressures the habitat may face; water quality parameters; levels of exposure to particular pressure, and more integrated indices which describe habitat structure and function, such as trophic index, or successional stages of development in habitats that have a natural cycle of change over time.

There are no commonly agreed indicators of quality for this habitat, although particular parameters may have been set in certain situations e.g. protected features within Natura 2000 sites, where reference values have been determined and applied on a location-specific basis. Examples of indicators of "naturalness" that are potential indicators for quality identified for offshore sands are; typical populations of bivalves and epifaunal brittlestars; maintained presence of substratum; lack of smothering; typically diverse communities with no increase in hardy or opportunistic species; and maintenance of sediment characteristics with typical levels of diversity.

Characteristic species:

Echinocardium cordatum and in deeper water the pea urchin *Echinocyamus pusillus*, the brittlestar *Amphiura filiformis*, *Ophiura albida* and *O. ophiura*, the anemone *Cerianthus lloydii*, and polychaete worms such as *Lanice conchilega*. In some locations the community is characterised by the bivalve *Abra prismatica*, the amphipod *Bathyporeia elegans* and polychaetes such as *Scoloplos armiger*, *Spiophanes bombyx*, *Aonides paucibranchiata*, *Chaetozone setosa*, *Ophelia borealis* and *Nephtys longosetosa*. Off the coast of mainland Portugal characteristic species are the sea urchin *Echinocardium cordatum*, bivalve *Macra stultorum*, polychaetes *Magelona johnstoni*, *Mediomastus fragilis*, *Owenia fusiformis* and *Spiophanes bombyx*. More mobile species reported from this habitat are *Pagurus bernhardus* and *Asterias rubens*.

Classification

EUNIS (v1405):

Level 4. A sub-habitat of 'Atlantic circalittoral sand' (A5.2).

Annex 1:

1110 Sandbanks slightly covered all the time

1160 Large shallow inlets and bays

MAES:

Marine - Marine inlets and transitional waters

Marine - Coastal

MSFD:

Shallow sublittoral sand

EUSeaMap:

Shallow sands

IUCN:

9.4 Subtidal sandy

Does the habitat type present an outstanding example of typical characteristics of one or more biogeographic regions?

Yes

Regions

Atlantic

Justification

This habitat is widespread and common in the North East Atlantic region.

Geographic occurrence and trends


Region	Present or Presence Uncertain	Current area of habitat	Recent trend in quantity (last 50 yrs)	Recent trend in quality (last 50 yrs)
<i>North-East Atlantic</i>	Bay of Biscay and the Iberian Coast: Present Celtic Seas: Present Greater North Sea: Present Macaronesia: Present Kattegat: Present	Unknown Km ²	Unknown	Decreasing

Extent of Occurrence, Area of Occupancy and habitat area

	Extent of Occurrence (EOO)	Area of Occupancy (AOO)	Current estimated Total Area	Comment
<i>EU 28</i>	4,005,871 Km ²	1,236	>10,997 Km ²	The area estimate for this habitat has been derived from a synthesis of EUNIS seabed habitat geospatial information for the European Seas but is recognised as being an underestimate.

	Extent of Occurrence (EOO)	Area of Occupancy (AOO)	Current estimated Total Area	Comment
EU 28+	>4,005,871 Km ²	>1,236	>10,997 Km ²	EOO and AOO have been calculated on the available data. Although this data set is known to be incomplete the figures exceed the thresholds for threatened status.

Distribution map



There are insufficient data to provide a comprehensive and accurate map of the distribution of this habitat. This map has been generated using EMODnet data from modelled/surveyed records for the North East Atlantic (and supplemented with expert opinion where applicable) (EMODnet 2010). EOO and AOO have been calculated on the available data presented in this map however these should be treated with caution as expert opinion is that this is not the full distribution of the habitat.

How much of the current distribution of the habitat type lies within the EU 28?

This habitat occurs in the EU 28+ (e.g. Norway, Isle of Man, Channel Islands). Percentage hosted by EU 28 is therefore less than 100% but there is insufficient information to establish the proportion.

Trends in quantity

It is difficult to establish the quantity of this habitat as it often has a patchy distribution, grading into other soft sediment habitats, or interspersed amongst rocky areas. Even where the extent of this habitat or its associated biotopes has been mapped in detail (e.g. as part of resource assessments for sand and gravel extraction or within marine protected areas) there is a lack of information on trends.

- Average current trend in quantity (extent)

EU 28: Unknown

EU 28+: Unknown

- Does the habitat type have a small natural range following regression?

No

Justification

This habitat has a widespread distribution in the North East Atlantic with the EOO >50,000km².

- Does the habitat have a small natural range by reason of its intrinsically restricted area?

No

Justification

This habitat has a widespread distribution in the North East Atlantic with the EOO >50,000km².

Trends in quality

Most sedimentary benthic systems on the continental shelf of Europe have been modified by fishing activities in the last 100 years, particularly by bottom trawls and dredging. Scientific evidence, supplemented with expert judgement to develop fisheries measures in protected areas for the Dutch sector of the North Sea, for example indicated that the most significant threat to the conservation status of the Dogger Bank comes from bottom gear, notably from beam trawling with tickler chains. The main effect

is on abiotic conditions, hence on structure and function, which results in reduction of the abundance of typical species. This initial effect is greater in sandy than muddy bottom however this is compensated somewhat by shorter recovery times where the seabed is predominantly sandy.

An analysis of the fishing intensity of EU trawlers (bottom otter, beam and mid-water trawls) using Automatic Identification System (AIS) ship tracking data over one year (2013/2014) shows high coverage in all European coastal waters and over the continental shelf. When combined with the modelled distribution of EUNIS marine habitat types it is possible to examine the extent of likely impact on a particular benthic habitat. For example, over this time period more than 40% of circalittoral fine sands and muddy sand were subject to trawling fishing pressure in the North Sea, with over 10% of this being interpreted a high or moderate pressure. When combining data for the North Sea and Celtic Sea more than 80% of this habitat type is considered to have been subject to such fishing pressure. Given that this is based on a single year of data and that this type of pressure has been taking place for decades it is likely to be an underestimate of the total area of this habitat affected by mobile demersal fishing gears.

- Average current trend in quality

EU 28: Decreasing

EU 28+: Decreasing

Pressures and threats

Disturbance of the substratum due to intensive fishing activities using bottom trawls or dredges may damage or modify infaunal communities, with burrowing echinoderms and bivalves being particularly vulnerable. Loss of substrate is likely to be detrimental to some biotopes where the majority of the characterising species are interstitial polychaetes. Smothering is not likely to cause considerable mortality to the species associated with the biotopes comprising this habitat type as most are able to reposition themselves at their preferred depth relatively quickly or are infaunal species. Polluted sediments may however cause a change in the species composition through changes in the sedimentary environment. Inshore dredging or construction works can cause changes to sediments through altered hydrodynamics or sedimentary processes, resulting in an altered community structure. In addition, synthetic and non-synthetic compound contamination from dumped dredge spoil and local discharges is also a possible threat that could cause a decline in the species richness within this habitat.

Brittlestars associated with this habitat, *O. fusiformis* and *A. filiformis* are likely to benefit from the

increased plankton production that occurs in areas of increased nutrient concentration, however, if enrichment leads to anoxia then reductions in growth rate and mass mortality may occur.

List of pressures and threats

Biological resource use other than agriculture & forestry

- Fishing and harvesting aquatic resources
 - Professional active fishing
 - Benthic or demersal trawling
 - Benthic dredging

Pollution

- Marine water pollution
 - Toxic chemical discharge from material dumped at sea
 - Non-synthetic compound contamination
 - Synthetic compound contamination

Natural System modifications

- Human induced changes in hydraulic conditions
 - Removal of sediments (mud...)
 - Estuarine and coastal dredging
 - Extraction of sea-floor and subsoil minerals (e.g. sand, gravel, rock, oil, gas)
 - Change of sea-floor substrate
 - Dykes, embankments, artificial beaches, general
 - Sea defense or coast protection works, tidal barrages
 - Dykes and flooding defense in inland water systems

Conservation and management

This habitat is afforded some protection within Marine Protected Areas. Beneficial management measures for this habitat include control of any kind of activity which damages or disturb seabed communities such as demersal fishing, dredging and offshore construction works. The regulation of effluent discharges can also support the conservation of this habitat.

List of conservation and management needs

Measures related to wetland, freshwater and coastal habitats

- Restoring/Improving water quality

Measures related to marine habitats

- Other marine-related measures

Measures related to spatial planning

- Establish protected areas/sites

Measures related to hunting, taking and fishing and species management

- Regulation/Management of fishery in marine and brackish systems

Conservation status

Annex 1:

1110: MATL U2, MMAC U1

1160: MATL U2, MMAC FV

When severely damaged, does the habitat retain the capacity to recover its typical character and functionality?

Large bodied, slow growing fauna such as bivalves which are associated with this habitat are sensitive to fishing disturbances and their populations may be slow to recover. Areas that are heavily fished may never fully recover because the seabed is re-disturbed before recovery has taken place. The timescales for recovery will depend on the individual area and the community present, bivalves of the genus *Thyasira* occur in isolated populations, and due to the lack of a dispersing larval stage, and are unlikely to recover if lost.

Thyasirids, small burrowing bivalves which live in fine sediments, are thought to be fairly slow growing and recovery of a damaged population is likely to take up to 5 years and depends on direct recruitment from the same population due to the low dispersal potential of these species. Where they occur in isolated populations they are therefore unlikely to recover if lost. In comparison, the high fecundity and larval dispersal potential of many of the polychaetes associated with this habitat is likely to result in a population recovering quite quickly - in less than a year for *O. fusiformis* and in approximately 5-6 years for *A. filiformis* due to the later age at which it reaches sexual maturity

Effort required

10 years
Naturally

Red List Assessment

Criterion A: Reduction in quantity

Criterion A	A1	A2a	A2b	A3
EU 28	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %

There are some estimates of the area of this habitat but insufficient information to determine overall trends in the quantity. It has therefore been assessed as Data Deficient under criterion A for both the EU 28 and EU 28+.

Criterion B: Restricted geographic distribution

Criterion B	B1				B2				B3
	EOO	a	b	c	AOO	a	b	c	
EU 28	>50,000 Km ²	Yes	Yes	No	>50	Yes	Yes	No	No
EU 28+	>50,000 Km ²	Yes	Yes	No	>50	Yes	Yes	No	No

This habitat has a large natural range in the North East Atlantic region. The precise extent is unknown however as EOO >50,000 km² and AOO >50, this exceeds the thresholds for a threatened category on the basis of restricted geographic distribution. There has been a decline in the biotic quality of this habitat and the major threat (mobile demersal fisheries) is likely to cause continuing declines in quality within the next 20 years, however the distribution of the habitat is such that the identified threats are unlikely to affect all localities at once. This habitat has therefore been assessed as Least Concern under criterion B.

Criterion C and D: Reduction in abiotic and/or biotic quality

Criteria C/D	C/D1		C/D2		C/D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	>80 %	Intermediate %	unknown %	unknown %	unknown %	unknown %
EU 28+	>80 %	Intermediate %	unknown %	unknown %	unknown %	unknown %

Criterion C	C1		C2		C3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %
EU 28+	unknown %	unknown %	unknown %	unknown %	unknown %	unknown %

Criterion D	D1		D2		D3	
	Extent affected	Relative severity	Extent affected	Relative severity	Extent affected	Relative severity
EU 28	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%
EU 28+	unknown %	unknown%	unknown %	unknown%	unknown %	unknown%

Substantial reductions in quality in at least some parts of this habitat are known to have occurred and continue to take place (as revealed by fishing distribution and intensity maps).

An analysis of the fishing intensity of EU trawlers (bottom otter, beam and mid-water trawls) using Automatic Identification System (AIS) ship tracking data over one year (2013/2014) shows high coverage in all European coastal waters and over the continental shelf. When combined with the modelled distribution of EUNIS marine habitat types it is possible to examine the extent of likely impact on a particular benthic habitat. For example, over this time period more than 40% of circalittoral fine sands and muddy sand were subject to trawling fishing pressure in the North Sea, with over 10% of this being interpreted a high or moderate pressure. When combining data for the North Sea and Celtic Sea more than 80% of this habitat type is considered to have been subject to such fishing pressure. Given that this is based on a single year of data, and that this type of pressure has been taking place for decades, it is likely to be an underestimate of the total area of this habitat which has been subject to such pressure.

Expert opinion is that there is likely to have been a very substantial reduction in quality of this habitat - an intermediate decline in quality affecting more than 80% of this habitat in the North East Atlantic region although it is also possible that more than 30% has been subject to a severe decline. This will depend on factors such as the intensity and frequency of disturbance. This habitat has therefore been assessed as Endangered under criteria C/D for both the EU 28 and EU 28+.

Criterion E: Quantitative analysis to evaluate risk of habitat collapse

Criterion E	Probability of collapse
EU 28	unknown
EU 28+	unknown

There is no quantitative analysis available to estimate the probability of collapse of this habitat type.

Overall assessment "Balance sheet" for EU 28 and EU 28+

	A1	A2a	A2b	A3	B1	B2	B3	C/D1	C/D2	C/D3	C1	C2	C3	D1	D2	D3	E
EU28	DD	DD	DD	DD	LC	LC	LC	EN	DD	DD	DD	DD	DD	DD	DD	DD	DD

	A1	A2a	A2b	A3	B1	B2	B3	C/D1	C/D2	C/D3	C1	C2	C3	D1	D2	D3	E
EU28+	DD	DD	DD	DD	LC	LC	LC	EN	DD	DD	DD	DD	DD	DD	DD	DD	DD

Overall Category & Criteria			
EU 28		EU 28+	
Red List Category	Red List Criteria	Red List Category	Red List Criteria
Endangered	C/D1	Endangered	C/D1

Confidence in the assessment

Medium (evenly split between quantitative data/literature and uncertain data sources and assured expert knowledge)

Assessors

North East Atlantic Working Group: S. Gubbay, G. Saunders, H. Tyler-Walters, N. Dankers, F. Otero-Ferrer, J. Forde, K. Fürhaupter, R. Haroun, N. Sanders.

Contributors

C. Karamita and the North East Atlantic Working Group: S. Gubbay, G. Saunders, H. Tyler-Walters, N. Dankers, F. Otero-Ferrer, J. Forde, K. Fürhaupter, R. Haroun, N. Sanders

Reviewers

A.Darr.

Date of assessment

14/12/2015

Date of review

19/01/2016

References

Connor, D.W., Allen, J.H., Golding, N. *et al.* 2004. The Marine Habitat Classification for Britain and Ireland Version 04.05 JNCC. [online] Peterborough: ISBN 1 861 07561 8. Available at: http://jncc.defra.gov.uk/pdf/04_05_introduction.pdf. (Accessed: 30/08/2014).

Eigaard Ole R. *et al.* "Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions" ICES J. Mar. Sci., doi = 10.1093/icesjms/fsv099.

Eleftheriou, A. and Basford, D. J. 1989. The macrobenthic infauna of the offshore northern North Sea. *Journal of the Marine Biological Association of the U.K* 69(1): 123-143.

European Environment Agency 2014. EUNIS habitat type hierarchical view. Available at: <http://eunis.eea.europa.eu/habitats-code-browser.jsp>. (Accessed: 22/08/2014).

Hill, J. & Wilson, E. 2008. *Amphiura filiformis*. A brittlestar. Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme. [online] Plymouth: Marine Biological Association of the United Kingdom. Available at: <http://www.marlin.ac.uk/speciesbenchmarks.php?speciesID=2503>. (Accessed 22/08/2014).

ICES. 2011. Fisheries Measures in Protected Areas (FIMPAS) within the Exclusive Economic Zone (EEZ) of the Dutch part of the North Sea: areas outside the 12 nautical miles zone: Proposals, Dates, Venue. FIMPAS Steering Group. 35 pp.

Lancaster, J. (Ed.), McCallum, S., Lowe A. C., Taylor, E., Chapman A. and Pomfret, J. 2014. *Development of*

detailed ecological guidance to support the application of the Scottish MPA selection guidelines in Scotland's seas. Inverness: Scottish Natural Heritage Commissioned Report No.491. Offshore Subtidal Sands and Gravels – supplementary document. Available at: <http://www.snh.gov.uk/docs/A1211261.pdf>. (Accessed: 10/10/2014).

Neal, K. & Avant, P. 2008. *Owenia fusiformis*. A tubeworm. *Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme*. [online] Plymouth: Marine Biological Association of the United Kingdom. Available at: <http://www.marlin.ac.uk/speciesbenchmarks.php?speciesID=4001>. (Accessed: 10/10/2015).

Nilsson, P. & Ziegler, F. Spatial distribution of fishing effort in relation to seafloor habitats in the Kattegat, a GIS analysis. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 17(4):421-440.

Rayment, W. J. 2008. *Venerid bivalves in circalittoral coarse sand or gravel*. *Marine Life Information Network: Biology and Sensitivity Key Information Sub-programme*. Plymouth: Marine Biological Association of the United Kingdom. Available at: <http://www.marlin.ac.uk/habitatsensitivity.php?habitatid=63&code=1997>. (Accessed: 30/08/2014).

Tempera, F., E. Atchoi, P. Amorim, J. Gomes-Pereira and Gonçalves, J. 2013. *Atlantic Area Marine Habitats. Adding new Macaronesian habitat types from the Azores to the EUNIS Habitat Classification*. Technica.