


Species name: **Coregonus oxyrinchus**
 Annex: **II*, IV**

Species group: **Fish**
 Regions: **ATL**


Assessments of conservation status at the European level (all biogeographical regions - EU25)



MS	Region	Conservation status assessment					Population size & unit	Population Trend
		Range	Population	Habitat	Future prospects	Overall		
EU25	ATL						7000 indiv.	-

ATL: The houting is an anadromous whitefish which spawns in large rivers from which the young migrate to the sea to develop and grow to maturity. They then return to their natal rivers to breed. The original distribution of the houting was in the Baltic Sea and eastern parts of the North Sea and several large rivers (eg Elbe, Rhine, Weser) into which adults migrated to spawn. The stocks of this species have greatly declined during the second half of the 20th century and several populations have disappeared. Species is recorded from Atlantic biogeographical region only from northern DK. Its overall assessment is "bad" because species was evaluated in 3 parameters as "bad".

**Assessments of conservation status as reported by Member states
(all biogeographical regions - EU25)**



MS	Region	Conservation status assessment					Size&unit	Population trend	Data quality
		Range	Population	Habitat	Future prospects	Overall			
DK	ATL						7000 - 7000 indiv.	-	2

Data quality is based on an assessment by each Member State, 1 = good, 2 = medium, 3 = poor

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2007 and covering the period 2001-2006. More detailed information is available at <http://biodiversity.eionet.europa.eu/article17>