


Species name: **Dioszeghyana schmidtii**
Annex: **II, IV**

Species group: **Invertebrates**
Regions: **MED PAN**

Assessments of conservation status at the European level (all biogeographical regions - EU25)




MS	Region	Conservation status assessment					Population size & unit	Population Trend
		Range	Population	Habitat	Future prospects	Overall		
EU25	MED							X
EU25	PAN						58 - 69 loc.	

A Ponto-Mediterranean species of moths. It occurs in the South-Eastern Europe and in the Black sea region. Within the EU it is known as well from Bulgaria and Romania. Its life cycle is linked to oak or *Acer tataricum*, so it can be found in various types of xerophilous oak forests.

The conservation status in the Pannonian biogeographical region is 'unfavourable inadequate'. The range and the population size are not sufficient to ensure a long-term viability. Suitable habitats occur usually as isolated fragments, which contributes to the sensitivity of the local populations. The status in the Mediterranean biogeographical region is 'unknown', better information is required.

Assessments of conservation status as reported by Member states (all biogeographical regions - EU25)



MS	Region	Conservation status assessment					Size&unit	Population trend	Data quality
		Range	Population	Habitat	Future prospects	Overall			
EL	MED						3 - (3) x	X	3
HU	PAN						50 - 60 loc.	+	3
SK	PAN						8 - 9 loc.	=	1

Data quality is based on as assessment by each Member State, 1 = good, 2 = medium, 3 = poor

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2007 and covering the period 2001-2006. More detailed information is available at <http://biodiversity.eionet.europa.eu/article17>