


Species name: **Gortyna borelii lunata**  
 Annex: **II, IV**

Species group: **Invertebrates**  
 Regions: **ATL CON MED PAN**

## Assessments of conservation status at the European level (all biogeographical regions - EU25)




MS	Region	Conservation status assessment					Population size & unit	Population Trend
		Range	Population	Habitat	Future prospects	Overall		
EU25	ATL						159 grids	-
EU25	CON							X
EU25	MED							X
EU25	PAN						121 loc.	+

The fisher's estuarine moth occurs in coastal areas, nearby sea-walls and tall-grass meadows with fluctuating humidity. Its distribution is limited by the limited distribution of its host plants (*Peucedanum* spp.).

The overall status of this species was different in all biogeographical regions. The assessments were as follows: Atlantic: unfavourable 'bad'; Continental: 'unknown'; Mediterranean: 'favourable'; and Pannonian: unfavourable 'inadequate'. Data from Germany are lacking. In the UK this species is threatened by sea-level rise and inappropriate land management. It is now protected in the UK. This species is considered to be rare everywhere.

## Assessments of conservation status as reported by Member states (all biogeographical regions - EU25)



MS	Region	Conservation status assessment					Size&unit	Population trend	Data quality
		Range	Population	Habitat	Future prospects	Overall			
FR	ATL						N/A x	-	
UK	ATL						6 - 6 loc.	=	1
DE	CON						11 - (11) x	X	1
FR	MED						N/A x	X	3
HU	PAN						121 - 121 loc.	+	2

Data quality is based on as assessment by each Member State, 1 = good, 2 = medium, 3 = poor

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2007 and covering the period 2001-2006. More detailed information is available at <http://biodiversity.eionet.europa.eu/article17>