MS Region

Annex: IV

Species name: Saga pedo

Species group: Invertebrates Regions: ALP ATL CON MED PAN

Population

Assessments of conservation status at the European level (all biogeographical regions - EU25)

| | Region | Range | Population | Habitat | prospects | Overall | size & unit | Trend |
|------|--------|-------|------------|---------|-----------|---------|---------------|-------|
| EU25 | ALP | | | | | | 29 - 50 loc. | - |
| EU25 | ATL | | | | | | | Х |
| EU25 | CON | | | | | | 47 grids | |
| EU25 | MED | | | | | | 64 grids | |
| EU25 | PAN | | | | | | 62 - 120 loc. | |
| | | | | | | | | |


Euturo

Conservation status assessment

The predatory bush cricket is an insect species which is present in southern Central Europe and large parts of Central Asia. However, in Europe its distribution is scattered.


The overall assessment for the different geographical regions differs very much. Only in the Atlantic region its status is assessed as 'favourable. In the Alpine and Continental region its status is 'inadequate', in the Mediterranean region it is 'bad' and in the Pannonian it is assessed as 'unknown'. However, in the Pannonian region (mainly Hungary) its habitat range and population are assessed as 'favourable'. More data from particularly Slovenia and Hungary are desired in order to improve the overall assessment. The 1996 IUCN conservation status currently is 'Vulnerable', but needs to be updated.

Population





Assessments of conservation status as reported by Member states (all biogeographical regions - EU25)



| MS | Region | | Conservatior | | | Population | Data | | |
|----|--------|-------|--------------|---------|---------------------|------------|---------------|---|---------|
| | | Range | Population | Habitat | Future prospects | Overall | Size&unit | | quality |
| AT | ALP | | | | | | 12 - 16 loc. | = | 1 |
| IT | ALP | | | | | | 7 - 9 loc. | - | 2 |
| SK | ALP | | | | | | 10 - 25 loc. | _ | 2 |
| FR | ATL | | | | | | N/A x | Х | |
| AT | CON | | | | | | 14 - 18 loc. | = | 1 |
| FR | CON | | | | | | N/A x | Х | |
| IT | CON | | | | | | 8 - 18 loc. | - | 2 |
| SI | CON | | | | | | N/A x | Х | 3 |
| ES | MED | | | | | | 6 - (6) loc. | Х | 3 |
| FR | MED | | | | | | N/A x | Х | |
| IT | MED | | | | | | 21 - 33 loc. | _ | 2 |
| CZ | PAN | | | | | | 10 - 10 loc. | + | 2 |
| ΗU | PAN | | | | | | 50 - 100 loc. | Х | 2 |
| SK | PAN | | | | | | 2 - 10 loc. | _ | 2 |

Data quality is based on as assessment by each Member State, 1 = good, 2 = medium, 3 = poor

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2007 and covering the period 2001-2006. More detailed information is available at http://biodiversity.eionet.europa.eu/article17