



Hamatocaulis lapponicus

Annex	II
Priority	No
Species group	Non-vascular plants
Regions	Alpine, Boreal

Hamatocaulis lapponicus grows in wet, mesotrophic and often spring-influenced mires or on lake shores. It is robust pleurocarp, mostly sparsely and irregularly branched, growing in glossy, green, purple, reddish-brown or blackish tufts. This is an eastern subarctic species and it is reported for Boreal region by Finland, Latvia and Sweden. It is also found in one place in Estonia (Vellak 2007) and assessed as Data Deficient (DD) in year 2007.

Overall regional assessment in the Boreal region is "Unfavourable Bad" due to all such parameters in Finland (78.6% of the species distribution). Swedish report also "Unfavourable Bad" due to the status of the range and population. Latvian population are comparatively small. It is also reported for Alpine region of Sweden. The assessment is "Unfavourable Inadequate" due to small range and population size. Trend is negative in the Boreal and stable in the Alpine region.

The threats to this species are the drainage of fens.


Changes in overall conservation status between 2001-06 and 2007-12 report are mostly caused by different methodical approach and better data rather than real change in conservation status in Boreal region. No changes in overall conservation status between 2001-06 and 2007-12 reports in Alpine region.

Better data is required from Sweden.

Species: *Hamatocaulis lapponicus*

Report under the Article 17 of the Habitats Directive

Assessment of conservation status at the European biogeographical level



Region	Conservation status (CS) of parameters				Current CS	Trend in CS	% in region	Previous CS	Reason for change
	Range	Population	Habitat	Future prospects					
ALP	U1	U1	FV	U1	U1	=	3	U1	
BOR	U2	U2	U1	U2	U2	-	97	U1	Not genuine

See the endnote for more informationⁱ

Species: *Hamatocaulis lapponicus*

Report under the Article 17 of the Habitats Directive

Assessment of conservation status at the Member State level



Hamatocaulis lapponicus

Distribution and conservation status at the Member State level

- | | |
|---|--|
|  Favourable |  EU Member States |
|  Unfavourable - inadequate |  Outside data coverage |
|  Unfavourable - bad |  Biogeographical region |
|  Unknown | |

The map shows both Conservation Status and distribution using a 10 km x 10 km grid. Conservation status is assessed at biogeographical level. Therefore the representation in each grid cell is only illustrative.

Species: *Hamatocaulis lapponicus*

Report under the Article 17 of the Habitats Directive

MS Region	Conservation status of parameters				Current CS	Trend in CS	% in region	Previous CS	Reason for change
	Range	Population	Habitat	Future prospects					
SE ALP	U1	U1	FV	U1	U1	=	100.0		
FI BOR	U1	U2	U1	U2	U2	=	73.3	U1	Better data
LV BOR	U1	U1	U1	U1	U1	=	16.7	U1	
SE BOR	U2	U2	XX	U1	U2	-	10.0	U2	

Knowing that not all changes in conservation status between the reporting periods were genuine, Member States were asked to give the reasons for changes in conservation status. Bulgaria and Romania only joined the EU in 2007 and Greece did not report for 2007-12 so no reason is given for change for these countries. Greek data shown above is from 2001-06.

Main pressures and threats reported by Member States

Member States were asked to report the 20 most important threats and pressures using an agreed hierarchical list which can be found on the [Article 17 Reference Portal](#). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. Pressures and threats were ranked in three classes 'high, medium and low importance'; the tables below only show threats and pressures classed as 'high', for some species there were less than ten threats or pressures reported as highly important.

Ten most frequently reported 'highly important' pressures

Code	Activity	Frequency
D05	Improved access to site	33
H01	Pollution to surface waters	33
H05	Soil pollution and solid waste (excl. discharges)	33

Ten most frequently reported 'highly important' threats

Code	Activity	Frequency
J02	Changes in water bodies conditions	50
K02	Vegetation succession/Biocenotic evolution	50

Species: *Hamatocaulis lapponicus*

Report under the Article 17 of the Habitats Directive

Proportion of population covered by the Natura 2000 network

For species listed in the Annex II of the Directive Member States were asked to report the population size within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size in the biogeographical/marine region.

Percentage of coverage by Natura 2000 sites in biogeographical/marine region

	ALP	BOR
FI		80
LV		100
SE	100	71

See the endnotes for more informationⁱⁱ

Most frequently reported conservation measures

For species listed in the Annex II of the Directive Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 17 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.1	Establish protected areas/sites	50
6.3	Legal protection of habitats and species	33
6.2	Establishing wilderness areas/ allowing succession	17

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2013 and covering the period 2007-2012. More detailed information, including the MS reports, is available at:

<http://bd.eionet.europa.eu/article17/reports2012/species/summary/?group=Non-vascular+plants&period=3&subject=Hamatocaulis+lapponicus>

Species: *Hamatocaulis lapponicus*

Report under the Article 17 of the Habitats Directive

i Assessment of conservation status at the European biogeographical level: Current Conservation Status (Current CS) shows the status for the reporting period 2007-2012, Previous Conservation Status (Previous CS) for the reporting period 2000-2006. Reason for change in conservation status between the reporting periods indicates whether the changes in the status were genuine or not genuine. Previous Conservation Status was not assessed for Steppic, Black Sea and Marine Black Sea regions. For these regions the Previous status is therefore considered as 'unknown'. The percentage of the species population occurring within the biogeographical/marine region (% in region) is calculated based on the area of GIS distribution.

ii Percentage of coverage by Natura 2000 sites in biogeographical/marine region: In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece. The values are only provided for regions, in which the occurrence of the species has been reported by the Member States.