



## Erodium rupicola

---

<b>Annex</b>	II, IV
<b>Priority</b>	Yes
<b>Species group</b>	Vascular plants
<b>Regions</b>	Mediterranean

The plant *Erodium rupicola* is endemic to the southeast of the Iberian Peninsula, Spain (Mediterranean region). It grows within sub-rupicolous and, less commonly, rupicolous communities, in areas with a moderate degree of nitrification. In the Sierra Nevada and the Sierra de Filabres, it occurs within the associations of *Centrantho nevadensis-Sedetum brevifolii* and *Diantho lusitani-Antirrhinetum rupestris*. In the Sierra de Baza, within the associations of *Teucrio rotundifolii-Kerneretum boissieri*. The IUCN European Red List classifies the species as Vulnerable (VU). The total population size of this species is estimated as 5,000 – 10,000 individuals and the area of occupancy has been reported 1,300 km<sup>2</sup>.

Both the current and previous conservation status are "Unknown". The country report of Spain indicates "Unknown" future prospects and unknown trend.

The species is threatened mostly by reduction or loss of specific habitat features, damage by herbivores and non intensive grazing. Forestry activities not referred to above and competition are also regarded as threats.


No changes in overall conservation status between 2001-06 and 2007-12 reports.

Better data required from Spain.

# Species: *Erodium rupicola*

Report under the Article 17 of the Habitats Directive

## Assessment of conservation status at the European biogeographical level




Region	Conservation status (CS) of parameters				Current CS	Trend in CS	% in region	Previous CS	Reason for change
	Range	Population	Habitat	Future prospects					
MED	FV	XX	XX	XX	XX	x	100	XX	

See the endnote for more information<sup>i</sup>

# Species: *Erodium rupicola*

Report under the Article 17 of the Habitats Directive

## Assessment of conservation status at the Member State level



### *Erodium rupicola*

Distribution and conservation status at the Member State level

- |                           |                        |
|---------------------------|------------------------|
| Favourable                | EU Member States       |
| Unfavourable – inadequate | Outside data coverage  |
| Unfavourable – bad        | Biogeographical region |
| Unknown                   |                        |

The map shows both Conservation Status and distribution using a 10 km x 10 km grid. Conservation status is assessed at biogeographical level. Therefore the representation in each grid cell is only illustrative.

# Species: *Erodium rupicola*

Report under the Article 17 of the Habitats Directive

MS	Region	Conservation status of parameters				Current CS	Trend in CS	% in region	Previous CS	Reason for change
		Range	Population	Habitat	Future prospects					
ES	MED	FV	XX	XX	XX		100.0	XX		

Knowing that not all changes in conservation status between the reporting periods were genuine, Member States were asked to give the reasons for changes in conservation status. Bulgaria and Romania only joined the EU in 2007 and Greece did not report for 2007-12 so no reason is given for change for these countries. Greek data shown above is from 2001-06.

## Main pressures and threats reported by Member States

Member States were asked to report the 20 most important threats and pressures using an agreed hierarchical list which can be found on the [Article 17 Reference Portal](#). Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. Pressures and threats were ranked in three classes 'high, medium and low importance'; the tables below only show threats and pressures classed as 'high', for some species there were less than ten threats or pressures reported as highly important.

### Ten most frequently reported 'highly important' pressures

Code	Activity	Frequency
A04	Grazing by livestock	33
J03	Other changes to ecosystems	33
K04	Interspecific floral relations	33

### Ten most frequently reported 'highly important' threats

Code	Activity	Frequency
A04	Grazing by livestock	33
J03	Other changes to ecosystems	33
K04	Interspecific floral relations	33

## Proportion of population covered by the Natura 2000 network

For species listed in the Annex II of the Directive Member States were asked to report the population size within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size in the biogeographical/marine region.

### Percentage of coverage by Natura 2000 sites in biogeographical/marine region

MED	
ES	32

See the endnotes for more information<sup>ii</sup>

# Species: *Erodium rupicola*

Report under the Article 17 of the Habitats Directive

## Most frequently reported conservation measures

For species listed in the Annex II of the Directive Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 17 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

### Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.3	Legal protection of habitats and species	50
7.4	Specific single species or species group management measures	50

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2013 and covering the period 2007-2012. More detailed information, including the MS reports, is available at:  
<http://bd.eionet.europa.eu/article17/reports2012/species/summary/?group=Vascular+plants&period=3&subject=Erodium+rupicola>

# Species: *Erodium rupicola*

Report under the Article 17 of the Habitats Directive

**i Assessment of conservation status at the European biogeographical level:** Current Conservation Status (Current CS) shows the status for the reporting period 2007-2012, Previous Conservation Status (Previous CS) for the reporting period 2000-2006. Reason for change in conservation status between the reporting periods indicates whether the changes in the status were genuine or not genuine. Previous Conservation Status was not assessed for Steppic, Black Sea and Marine Black Sea regions. For these regions the Previous status is therefore considered as 'unknown'. The percentage of the species population occurring within the biogeographical/marine region (% in region) is calculated based on the area of GIS distribution.

**ii Percentage of coverage by Natura 2000 sites in biogeographical/marine region:** In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (\*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece. The values are only provided for regions, in which the occurrence of the species has been reported by the Member States.