# **European Environment Agency** *European Topic Centre on Biological Diversity*



## Ornithogalum reverchonii

Annex IV Priority No

**Species group** Vascular plants **Regions** Mediterranean

The plant *Ornithogalum reverchonii* is endemic to Spain and Morocco (Mediterranean region). This species is found in vertical limestone shadow cliffs, in perennial grasslands growing in rock crevices or ledges in peaty soils. The IUCN Red List classifies the species as Least Concern (LC). It is included as Vulnerable (VU) in the regional catalogue of threatened wild flora of Andalucía.

Little information provided by Spain, therefore the previous and current conservation status is "Unknown". Only range was reported as "Favourable".

There is no information on potential threats available.


No changes in overall conservation status between 2001-06 and 2007-12 reports.

Better data required from Spain.

# Species: Ornithogalum reverchonii


Report under the Article 17 of the Habitats Directive

# Assessment of conservation status at the European biogeographical level



See the endnote for more information<sup>i</sup>

#### Assessment of conservation status at the Member State level



The map shows both Conservation Status and distribution using a  $10 \text{ km} \times 10 \text{ km}$  grid. Conservation status is assessed at biogeographical level. Therefore the representation in each grid cell is only illustrative.

Unknown

# Species: Ornithogalum reverchonii

Report under the Article 17 of the Habitats Directive

MS Region	Conservation status of parameters				Current	Trend in	% in	Previous	Reason
	Range	Population	Habitat	Future prospects	CS	CS	region	CS	for change
ES MED	FV	XX	XX	XX	XX		100.0	XX	

Knowing that not all changes in conservation status between the reporting periods were genuine, Member States were asked to give the reasons for changes in conservation status. Bulgaria and Romania only joined the EU in 2007 and Greece did not report for 2007-12 so no reason is given for change for these countries. Greek data shown above is from 2001-06.

## Main pressures and threats reported by Member States

Member States were asked to report the 20 most important threats and pressures using an agreed hierarchical list which can be found on the Article 17 Reference Portal. Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. Pressures and threats were ranked in three classes 'high, medium and low importance'; the tables below only show threats and pressures classed as 'high', for some species there were less than ten threats or pressures reported as highly important.

Ten most frequently reported 'highly important' pressures

Code Activity	Frequency							
No 'highly important' pressures were reported.								
Ten most frequently reported 'highly important' threats								
Code Activity	Frequency							
No 'highly important' threats were reported.								

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2013 and covering the period 2007-2012. More detailed information, including the MS reports, is available at: http://bd.eionet.europa.eu/article17/reports2012/species/summary/? group=Vascular+plants&period=3&subject=Ornithogalum+reverchonii

# Species: Ornithogalum reverchonii

Report under the Article 17 of the Habitats Directive

Assessment of conservation status at the European biogeographical level: Current Conservation Status (Current CS) shows the status for the reporting period 2007-2012, Previous Conservation Status (Previous CS) for the reporting period 2000-2006. Reason for change in conservation status between the reporting periods indicates whether the changes in the status were genuine or not genuine. Previous Conservation Status was not assessed for Steppic, Black Sea and Marine Black Sea regions. For these regions the Previous status is therefore considered as 'unknown'. The percentage of the species population occurring within the biogeographical/marine region (% in region) is calculated based on the area of GIS distribution.